

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



UNITED STATES PATENT APPLICATION

OF

DAVID F. CARMICHAEL,

DAVID C. ANDERSON,

GEORGE P. STRICKLIN,

AND

HOWARD G. WELGUS

FOR

METALLOPROTEINASE INHIBITOR SEQUENCE,

RECOMBINANT VECTOR SYSTEM FOR USING SAME AND

RECOMBINANT-DNA METHOD FOR THE MANUFACTURE OF SAME

09987367-01303
205270-75E78660

Justin

BACKGROUND OF THE INVENTION

[001] This is a continuation-in-part application of U.S. Patent Application Serial No. 699,181, filed February 5, 1985. Endogenous proteolytic enzymes serve to degrade invading organisms, antigen-antibody complexes and certain tissue proteins which are no longer necessary or useful to the organism. In a normally functioning organism, proteolytic enzymes are produced in a limited quantity and are regulated in part through specific inhibitors.

[002] Metalloproteinases are enzymes present in the body which are often involved in the degradation of connective tissue. While some connective tissue degradation is necessary for normal functioning of an organism, an excess of connective tissue degradation occurs in several disease states and is believed to be attributable, at least in part, to excess metalloproteinase. It is believed that metalloproteinases are at least implicated in periodontal disease, corneal and skin ulcers, rheumatoid arthritis and the spread of cancerous solid tumors.

[003] These diseases generally occur in areas of the body which contain a high proportion of collagen, a particular form of connective tissue. An examination of patients with these diseases of connective tissue has revealed an excessive breakdown of the various components of connective tissues, including collagen proteoglycans and elastin. Therefore, it has been deduced that an excessive concentration of a particular metalloproteinase, for example collagenase, proteoglyconase, gelatinase, and certain elastases, may cause or exacerbate the connective tissue destruction associated with the aforementioned diseases.

[004] In the normal state, the body possesses metalloproteinase inhibitors which bind to metalloproteinases to effectively prevent these enzymes from acting on

their connective tissue substrates. Specifically, in a healthy organism, metalloproteinase inhibitors are present in concentrations sufficient to interact with metalloproteinases to an extent which allows sufficient quantities of metalloproteinase to remain active while binding the excess metalloproteinase so that the connective tissue damage seen in the various diseases does not occur.

[005] It is postulated that one immediate cause of the connective tissue destruction present in the foregoing disease states is an imbalance in the relative metalloproteinase/metalloproteinase inhibitor concentrations. In these situations, either due to an excessive amount of active metalloproteinase or a deficiency in the amount of active metalloproteinase inhibitor, the excess metalloproteinase is believed to cause the connective tissue degradation responsible for causing or exacerbating the disease. It is postulated that, by treating persons with connective tissue diseases with metalloproteinase inhibitors, the degradative action of the excess metalloproteinase may be curtailed or halted. Therefore, particular metalloproteinase inhibitors of specific interest to the present inventors are collagenase inhibitors because it is believed that these inhibitors would be pharmaceutically useful in the treatment or prevention of connective tissue diseases.

[006] The existence of metalloproteinase and metalloproteinase inhibitors has been discussed in the scientific literature. For example, Sellers *et al.*, Biochemical And Biophysical Research Communications 87:581-587 (1979), discusses isolation of rabbit bone collagenase inhibitor. Collagenase inhibitor isolated from human skin fibroblasts is discussed in Stricklin and Welgus, J.B.C. 258:12252-12258 (1983) and Welgus and Stricklin, J.B.C. 258:12259-12264 (1983). The presence of collagenase inhibitors in

naturally-occurring body fluids is further discussed in Murphy *et al.*, *Biochem. J.* 195:167-170 (1981) and Cawston *et al.*, *Arthritis and Rheumatism*, 27:285 (1984). In addition, metalloproteinase inhibitors are discussed by Reynolds *et al.* in Cellular Interactions, Dingle and Gordon, eds., (1981). Although these articles characterize particular, isolated metalloproteinase inhibitors and discuss, to some extent, the role or potential role of metalloproteinases in connective tissue disease treatment and speculate on the ability of metalloproteinase inhibitors to counteract this destruction, none of these researchers had previously been able to isolate a portable DNA sequence capable of directing intracellular production of metalloproteinase inhibitors or to create a recombinant-DNA method for the production of these inhibitors.

[007] Surprisingly, the present inventors have discovered a portable DNA sequence capable of directing the recombinant-DNA synthesis of metalloproteinase inhibitors. These metalloproteinase inhibitors are biologically equivalent to those isolated from human skin fibroblast cultures. The metalloproteinase inhibitors of the present invention, prepared by the recombinant-DNA methods set forth herein, will enable increased research into prevention and treatment of metalloproteinase-induced connective tissue diseases. In addition, the metalloproteinase inhibitors of the present invention are useful in neutralizing metalloproteinases, including the excess metalloproteinase associated with disease states. Therefore, it is believed that a cure for these diseases will be developed which will embody, as an active ingredient, the metalloproteinase inhibitors of the present invention. Furthermore, the metalloproteinase inhibitors of the present invention are capable of interacting with their

metalloproteinase targets in a manner which allows the development of diagnostic tests for degradative connective tissue diseases using the newly discovered inhibitors.

[008] The recombinant metalloproteinase inhibitors discussed herein interact stoichiometrically (*i.e.*, in a 1:1 ratio) with their metalloproteinase targets. In addition, these metalloproteinase inhibitors are heat resistant, acid stable, glycosylated, and exhibit a high isoelectric point.

SUMMARY OF THE INVENTION

[009] The present invention relates to metalloproteinase inhibitors and a recombinant-DNA method of producing the same and to portable DNA sequences capable of directing intracellular production of the metalloproteinase inhibitors. Particularly, the present invention relates to a collagenase inhibitor, a recombinant-DNA method for producing the same and to portable DNA sequences for use in the recombinant method. The present invention also relates to a series of vectors containing these portable DNA sequences.

[010] One object of the present invention is to provide a metalloproteinase inhibitor, which can be produced in sufficient quantities and purities to provide economical pharmaceutical compositions which possess metalloproteinase inhibitor activity.

[011] An additional object of the present invention is to provide a recombinant-DNA method for the production of these metalloproteinase inhibitors. The recombinant metalloproteinase inhibitors produced by this method are biologically equivalent to the metalloproteinase inhibitor isolable from human skin fibroblast cultures.

[012] To facilitate the recombinant-DNA synthesis of these metalloproteinase inhibitors, it is a further object of the present invention to provide portable DNA sequences capable of directing intracellular production of metalloproteinase inhibitors. It is also an object of the present invention to provide cloning vectors containing these portable sequences. These vectors are capable of being used in recombinant systems to produce pharmaceutically useful quantities of metalloproteinase inhibitors.

[013] Additional objects and advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description or may be learned from practice of the invention. The objects and advantages may be realized and attained by means of the instrumentalities and combinations particularly pointed out in the appended claims.

[014] To achieve the objects and in accordance with the purposes of the present invention, metalloproteinase inhibitors are set forth, which are capable of stoichiometric reaction with metalloproteinases. These metalloproteinase inhibitors are remarkably heat resistant, acid stable, glycosylated, and exhibit a high isoelectric point. Furthermore, these metalloproteinase inhibitors are biologically equivalent to those inhibitors isolated from human skin fibroblast cultures.

[015] To further achieve the objects and in accordance with the purposes of the present invention, as embodied and broadly described herein, portable DNA sequences coding for metalloproteinase inhibitors are provided. These sequences comprise nucleotide sequences capable of directing intracellular production of metalloproteinase inhibitors. The portable sequences may be either synthetic sequences or restriction fragments ("natural" DNA sequences). In a preferred

embodiment, a portable DNA sequence is isolated from a human fibroblast cDNA library and is capable of directing intracellular production of a collagenase inhibitor which is biologically equivalent to that inhibitor which is from a human skin fibroblast culture.

[016] The coding strand of a first preferred DNA sequence which has been discovered has the following nucleotide sequence:

10 20 30 40 50 60
GTTGTTGCTG TGGCTGATAG CCCAGCAGG GCCTGCACCT GTGTCCCACC CCACCCACAG

70 80 90 100 110 120
ACGGCCTTCT GCAATTCCGA CCTCGTCATC AGGGCCAAGT TCGTGGGGAC ACCAGAAGTC

130 140 150 160 170 180
AACCAGACCA CCTTATACCA GCGTTATGAG ATCAAGATGA CCAAGATGTA TAAAGGGTTC

190 200 210 220 230 240
CAAGCCTTAG GGGATGCCGC TGACATCCGG TTCGTCTACA CCCCCGCCAT GGAGAGTGTC

250 260 270 280 290 300
TGCGGATACT TCCACAGGTC CCACAACCGC AGCGAGGAGT TTCTCATTGC TGGAAAACTG

310 320 330 340 350 360
CAGGATGGAC TCTTGACAT CACTACCTGC AGTTTCGTGG CTCCCTGGAA CAGCCTGAGC

370 380 390 400 410 420
TTAGCTCAGC GCCGGGGCTT CACCAAGACC TACACTGTTG GCTGTGAGGA ATGCACAGTG

430 440 450 460 470 480
TTTCCCTGTT TATCCATCCC CTGCAAACCTG CAGAGTGGCA CTCATTGCTT GTGGACGGAC

490 500 510 520 530 540
CAGCTCCTCC AAGGCTCTGA AAAGGGCTTC CAGTCCCGTC ACCTTGCCCTG CCTGCCTCGG

550 560 570 580 590 600
GAGCCAGGGC TGTGCACCTG GCAGTCCCTG CGGTCCGAGA TAGCCTGAAT CCTGCCCCGGA

610 620 630 640 650 660
GTGGAAGCTG AAGCCTGCAC AGTGTCCACC CTGTTCCGAC TCCCATCTTT CTTCCGGACA

670 680 690 700
ATGAAATAAA GAGTTACCAC CCAGCAAAAA AAAAAAGGAA TTC

[017] The nucleotides represented by the foregoing abbreviations are set forth in the Detailed Description of the Preferred Embodiments.

[018] A second preferred DNA sequence has been discovered which has an additional nucleotide sequence 5' to the initiator sequence. This sequence, which contains as the eighty-second through four-hundred-thirty-second nucleotides nucleotides 1 through 351 of the first preferred sequence set forth above, has the following nucleotide sequence:

10 20 30 40 50 60
GGCCATCGCC GCAGATCCAG CGCCCAGAGA GACACCAGAG AACCCACCAT GGCCCCCTTT

70 80 90 100 110 120
GACCCCTGGC TTCTGCATCC TGTGTGTGCT GTGGCTGATA GCCCCAGCAG GGCCTGCACC

130 140 150 160 170 180
TGTGTCCAC CCCACCCACA GACGGCCTTC TGCAATTCCG ACCTCGTCAT CAGGGCCAAG

190 200 210 220 230 240
TTCGTGGGGA CACCAGAAGT CAACCAGACC ACCTTATACC AGCGTTATGA GATCAAGATG

250 260 270 280 290 300
ACCAAGATGT ATAAAGGGTT CCAAGCCTTA GGGGATGCCG CTGACATCCG GTTCGTCTAC

310 320 330 340 350 360
ACCCCCGCCA TGGAGAGTGT CTGCGGATAC TTCCACAGGT CCCACAACCG CAGCGAGGAG

370 380 390 400 410 420
TTTCTCATTG CTGGAAACT GCAGGATGGA CTCTTGACA TCACTACCTG CAGTTTCGTG

430
GCTCCCTGGA AC

[019] A third preferred DNA sequence which incorporates the 5' region of the second preferred sequence and the 3' sequence of the first preferred sequence, has the following nucleotide sequence:

10 20 30 40 50 60
GGCCATCGCC GCAGATCCAG CGCCCAGAGA GACACCAGAG AACCCACCAT GGCCCCCTTT

70 80 90 100 110 120

GACCCCTGGC TTCTGCATCC TGTGTTGCT GTGGCTGATA GCCCCAGCAG GGCCTGCACC
 130 140 150 160 170 180
 TGTGTCCCAC CCCACCCACA GACGGCCTTC TGCAATTCCG ACCTCGTCAT CAGGGCCAAG
 190 200 210 220 230 240
 TTCGTGGGGA CACCAGAAGT CAACCAGACC ACCTTATACC AGCGTTATGA GATCAAGATG
 250 260 270 280 290 300
 ACCAAGATGT ATAAAGGGTT CCAAGCCTTA GGGGATGCCG CTGACATCCG GTTCGTCTAC
 310 320 330 340 350 360
 ACCCCCGCCA TGGAGAGTGT CTGCGGATAC TTCCACAGGT CCCACAACCG CAGCGAGGAG
 370 380 390 400 410 420
 TTTCTCATTG CTGGAAACT GCAGGATGGA CTCTGCACA TCACTACCTG CAGTTTCGTG
 430 440 450 460 470 480
 TCTCCCTGGA ACAGCCTGAG CTTAGCTCAG CGCCGGGGCT TCACCAAGAC CTACACTGTT
 490 500 510 520 530 540
 GGCTGTGAGG AATGCACAGT GTTTCCCTGT TTATCCATCC CCTGCAAACCT GCAGAGTGGC
 550 560 570 580 590 600
 ACTCATTGCT TGTGGACGGA CCAGCTCCTC CAAGGCTCTG AAAAGGGCTT CCAGTCCCGT
 610 620 630 640 650 660
 CACCTTGCCT GCCTGCCTCG GGAGCCAGGG CTGTGCACCT GGCAGTCCCT GCGGTCCGAG
 670 680 690 700 710 720
 ATAGCCTGAA TCCTGCCCCG AGTGAAGCT GAAGCCTGCA CAGTGTCCAC CCTGTTCCCA
 730 740 750 760 770 780
 CTCCCATCTT TCTTCCGGAC AATGAAATAA AGAGTTACCA CCCAGCAAAA AAAAAAAGGA

[020] To facilitate identification and isolation of natural DNA sequences for use
 in the present invention, the inventors have developed a human skin fibroblast cDNA
 library. This library contains the genetic information capable of directing a cell to
 synthesize the metalloproteinase inhibitors of the present invention. Other natural DNA
 sequences which may be used in the recombinant DNA methods set forth herein may
 be isolated from human genomic libraries.

[021] Additionally, portable DNA sequences useful in the processes of the present invention may be synthetically created. These synthetic DNA sequences may be prepared by polynucleotide synthesis and sequencing techniques known to those of ordinary skill in the art.

[022] Additionally, to achieve the objects and in accordance with the purposes of the present invention, a recombinant-DNA method is disclosed which results in microbial manufacture of the instant metalloproteinase inhibitors using the portable DNA sequences referred to above. This recombinant DNA method comprises:

- (a) preparation of a portable DNA sequence capable of directing a host microorganism to produce a protein having metalloproteinase inhibitor activity, preferably collagenase inhibitor activity;
- (b) cloning the portable DNA sequence into a vector capable of being transferred into and replicating in a host microorganism, such vector containing operational elements for the portable DNA sequence;
- (c) transferring the vector containing the portable DNA sequence and operational elements into a host microorganism capable of expressing the metalloproteinase inhibitor protein;
- (d) culturing the host microorganism under conditions appropriate for amplification of the vector and expression of the inhibitor; and
- (e) in either order:
 - (i) harvesting the inhibitor; and
 - (ii) causing the inhibitor to assume an active, tertiary structure whereby it possesses metalloproteinase inhibitor activity.

[023] To further accomplish the objects and in further accord with the purposes of the present invention, a series of cloning vectors are provided comprising at least one of the portable DNA sequences discussed above. In particular, plasmid pUC9-F5/237P10 is disclosed.

[024] It is understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

[025] The accompanying drawing, which is incorporated in and constitutes a part of this specification illustrates one embodiment of the invention and, together with the description, serves to explain the principles of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[026] Figure 1 is a partial restriction map of the plasmid pUC9-F5/237P10.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[027] Reference will now be made in detail to the presently preferred embodiments of the invention, which, together with the drawing and the following examples, serve to explain the principles of the invention.

[028] As noted above, the present invention relates in part to portable DNA sequences capable of directing intracellular production of metalloproteinase inhibitors in a variety of host microorganisms. "Portable DNA sequence" in this context is intended to refer either to a synthetically-produced nucleotide sequence or to a restriction fragment of a naturally occurring DNA sequence. For purposes of this specification, "metalloproteinase inhibitor" is intended to mean the primary structure of the protein as defined by the codons present in the deoxyribonucleic acid sequence which directs intracellular production of the amino acid sequence, and which may or may not include

post-translational modifications. It is contemplated that such post-translational modifications include, for example, glycosylation. It is further intended that the term "metalloproteinase inhibitor" refer to either the form of the protein as would be excreted from a microorganism or the methionyl-metalloproteinase inhibitor as it may be present in microorganisms from which it was not excreted.

[029] In a preferred embodiment, the portable DNA sequences are capable of directing intracellular production of collagenase inhibitors. In a particularly preferred embodiment, the portable DNA sequences are capable of directing intracellular production of a collagenase inhibitor biologically equivalent to that previously isolated from human skin fibroblast cultures. By "biologically equivalent", as used herein in the specification and claims, it is meant that an inhibitor, produced using a portable DNA sequence of the present invention, is capable of preventing collagenase-induced tissue damage of the same type, but not necessarily to the same degree, as a native human collagenase inhibitor, specifically that native human collagenase inhibitor isolable from human skin fibroblast cell cultures.

[030] A first preferred portable DNA sequence of the present invention has a nucleotide sequence as follows:

10 20 30 40 50 60
GTTGTTGCTG TGGCTGATAG CCCCAGCAGG GCCTGCACCT GTGTCCCACC CCACCCACAG
70 80 90 100 110 120
ACGGCCTTCT GCAATTCCGA CCTCGTCATC AGGGCCAAGT TCGTGGGGAC ACCAGAAGTC
130 140 150 160 170 180
AACCAGACCA CCTTATACCA GCGTTATGAG ATCAAGATGA CCAAGATGTA TAAAGGGTTC
190 200 210 220 230 240
CAAGCCTTAG GGGATGCCGC TGACATCCGG TTCGTCTACA CCCCCGCCAT GGAGAGTGTC

250 260 270 280 290 300
 TCGGATACT TCCACAGGTC CCACAACCGC AGCGAGGAGT TTCTCATTGC TGGAAAAC TG
 310 320 330 340 350 360
 CAGGATGGAC TCTTGACAT CACTACCTGC AGTTTCGTGG CTCCCTGGAA CAGCCTGAGC
 370 380 390 400 410 420
 TTAGCTCAGC GCCGGGGCTT CACCAAGACC TACACTGTTG GCTGTGAGGA ATGCACAGTG
 430 440 450 460 470 480
 TTTCCCTGTT TATCCATCCC CTGCAAACCTG CAGAGTGGCA CTCATTGCTT GTGGACGGAC
 490 500 510 520 530 540
 CAGCTCCTCC AAGGCTCTGA AAAGGGCTTC CAGTCCCCTG ACCTTGCCCTG CCTGCCTCGG
 550 560 570 580 590 600
 GAGCCAGGGC TGTGCACCTG GCAGTCCCTG CGGTCCCAGA TAGCCTGAAT CCTGCCCCGA
 610 620 630 640 650 660
 GTGGAAGCTG AAGCCTGCAC AGTGTCCACC CTGTTCCAC TCCCATCTTT CTTCCGGACA
 670 680 690 700
 ATGAAATAAA GAGTTACCAC CCAGCAAAAA AAAAAAGGAA TTC

wherein the following nucleotides are represented by the abbreviations indicated below.

<u>Nucleotides</u>	<u>Abbreviation</u>
Deoxyadenylic acid	A
Deoxyguanylic acid	G
Deoxycytidylic acid	C
Thymidylic acid	T

INS (5) [031] A second preferred portable DNA sequence of the present invention has
 the following nucleotide sequence:

10 20 30 40 50 60
 GGCCATCGCC GCAGATCCAG CGCCCAGAGA GACACCAGAG AACCCACCAT GGCCCCCTTT
 70 80 90 100 110 120
 GACCCCTGGC TTCTGCATCC TGTGTGTTGCT GTGGCTGATA GCCCCAGCAG GGCCTGCACC
 130 140 150 160 170 180
 TGTGTCCAC CCCACCCACA GACGGCCTTC TGCAATTCCG ACCTCGTCAT CAGGGCCAAG

190 200 210 220 230 240
TTCGTGGGGA CACCAGAAGT CAACCAGACC ACCTTATACC AGCGTTATGA GATCAAGATG
250 260 270 280 290 300
ACCAAGATGT ATAAAGGGTT CCAAGCCTTA GGGGATGCCG CTGACATCCG GTTCGTCTAC
310 320 330 340 350 360
ACCCCCGCCA TGGAGAGTGT CTGCGGATAC TTCCACAGGT CCCACAACCG CAGCGAGGAG
370 380 390 400 410 420
TTTCTCATTG CTGGAAACT GCAGGATGGA CTCTTGCACA TCACTACCTG CAGTTTCGTG
430
GCTCCCTGGA AC

[032] In this second preferred sequence, an open reading frame exists from nucleotides 1 through 432. The first methionine of this reading frame is encoded by nucleotides 49 through 51 and is the site of translation initiation. It should be noted that the amino acid sequence prescribed by nucleotides 49 through 114 is not found in the mature metalloproteinase. It is believed that this sequence is the leader peptide of the human protein.

[033] A third preferred portable DNA sequence has the nucleotide sequence:

10 20 30 40 50 60
GGCCATCGCC GCAGATCCAG CGCCCAGAGA GACACCAGAG AACCCACCAT GGCCCCCTTT
70 80 90 100 110 120
GACCCCTGGC TTCTGCATCC TGTGTGTGCT GTGGCTGATA GCCCAGCAG GGCCTGCACC
130 140 150 160 170 180
TGTGTCCCAC CCCACCCACA GACGGCCTTC TGCAATTCCG ACCTCGTCAT CAGGGCCAAG
190 200 210 220 230 240
TTCGTGGGGA CACCAGAAGT CAACCAGACC ACCTTATACC AGCGTTATGA GATCAAGATG
250 260 270 280 290 300
ACCAAGATGT ATAAAGGGTT CCAAGCCTTA GGGGATGCCG CTGACATCCG GTTCGTCTAC
310 320 330 340 350 360
ACCCCCGCCA TGGAGAGTGT CTGCGGATAC TTCCACAGGT CCCACAACCG CAGCGAGGAG
370 380 390 400 410 420
TTTCTCATTG CTGGAAACT GCAGGATGGA CTCTTGCACA TCACTACCTG CAGTTTCGTG

430 440 450 460 470 480
 GCTCCCTGGA ACAGCCTGAG CTTAGCTCAG CGCCGGGGCT TCACCAAGAC CTACACTGTT
 490 500 510 520 530 540
 GGCTGTGAGG AATGCACAGT GTTTCCCTGT TTATCCATCC CCTGCAAACCT GCAGAGTGGC
 550 560 570 580 590 600
 ACTCATTGCT TGTGGACGGA CCAGCTCCTC CAAGGCTCTG AAAAGGGCTT CCAGTCCCGT
 610 620 630 640 650 660
 CACCTTGCTT GCCTGCCTCG GGAGCCAGGG CTGTGCACCT GGCAGTCCCT GCGGTCCCGA
 670 680 690 700 710 720
 ATAGCCTGAA TCCTGCCCCG AGTGGAAGCT GAAGCCTGCA CAGTGTCCAC CCTGTTCCCA
 730 740 750 760 770 780
 CTCCCATCTT TCTTCCGGAC AATGAAATAA AGAGTTACCA CCCAGCAAAA AAAAAAAGGA

SUBC7
 cont

[034] This third sequence contains the 5' nontranslated region of the second preferred sequence and the 3' region of the first preferred sequence. It is envisioned that this third preferred sequence is capable of directing intracellular production of a metalloproteinase analogous to a mature human collagenase inhibitor in a microbial or mammalian expression system.

[035] It must be borne in mind in the practice of the present invention that the alteration of some amino acids in a protein sequence may not affect the fundamental properties of the protein. Therefore, it is also contemplated that other portable DNA sequences, both those capable of directing intracellular production of identical amino acid sequences and those capable of directing intracellular production of analogous amino acid sequences which also possess metalloproteinase inhibitor activity, are included within the ambit of the present invention.

[036] It is contemplated that some of these analogous amino acid sequences will be substantially homologous to native human metalloproteinase inhibitors while other amino acid sequences, capable of functioning as metalloproteinase inhibitors, will

not exhibit substantial homology to native inhibitors. By "substantial homology", as used herein, is meant a degree of homology to a native metalloproteinase inhibitor in excess of 50%, preferably in excess of 60%, preferably in excess of 80%. The percentage homology as discussed herein is calculated as the percentage of amino acid residues found in the smaller of the two sequences that align with identical amino acid residues in the sequence being compared when four gaps in a length of 100 amino acids may be introduced to assist in that alignment as set forth by Dayhoff, M.O. in Atlas of Protein Sequence and Structure Vol. 5, p. 124 (1972), National Biochemical Research Foundation, Washington, D.C., specifically incorporated herein by reference.

[037] As noted above, the portable DNA sequences of the present invention may be synthetically created. It is believed that the means for synthetic creation of these polynucleotide sequences are generally known to one of ordinary skill in the art, particularly in light of the teachings contained herein. As an example of the current state of the art relating to polynucleotide synthesis, one is directed to Matteucci, M.D. and Caruthers, M.H., in J. Am. Chem. Soc. 103: 3185 (1981) and Beaucage, S.L. and Caruthers, M.H. in Tetrahedron Lett. 22: 1859 (1981), specifically incorporated herein by reference.

[038] Additionally, the portable DNA sequence may be a fragment of a natural sequence, *i.e.*, a fragment of a polynucleotide which occurred in nature and which has been isolated and purified for the first time by the present inventors. In one embodiment, the portable DNA sequence is a restriction fragment isolated from a cDNA library. In this preferred embodiment, the cDNA library is created from human skin fibroblasts.

[039] In an alternative embodiment, the portable DNA sequence is isolated from a human genomic library. An example of such a library useful in this embodiment is set forth in Lawn *et al.* Cell 15: 1157-1174 (1978), specifically incorporated herein by reference.

[040] As also noted above, the present invention relates to a series of vectors, each containing at least one of the portable DNA sequences described herein. It is contemplated that additional copies of the portable DNA sequence may be included in a single vector to increase a host microorganism's ability to produce large quantities of the desired metalloproteinase inhibitor.

[041] In addition, the cloning vectors within the scope of the present invention may contain supplemental nucleotide sequences preceding or subsequent to the portable DNA sequence. These supplemental sequences are those that will not interfere with transcription of the portable DNA sequence and will, in some instances as set forth more fully hereinbelow, enhance transcription, translation, or the ability of the primary amino acid structure of the resultant metalloproteinase inhibitor to assume an active, tertiary form.

[042] A preferred vector of the present invention is set forth in Figure 1. This vector, pUC9-F5/237P10, contains the preferred nucleotide sequence set forth above. Vector pUC9-F5/237P10 is present in the C600/pUC9-F5/237P10 cells on deposit in the American Type Culture Collection in Rockville, Maryland under Accession No. 53003.

[043] A preferred nucleotide sequence encoding the metalloproteinase inhibitor is identified in Figure 1 as region A. Plasmid pUC9-F5/237P10 also contains

supplemental nucleotide sequences preceding and subsequent to the preferred portable DNA sequence in region A. These supplemental sequences are identified as regions B and C, respectively.

[044] In alternate preferred embodiments, either one or both of the preceding or subsequent supplemental sequences may be removed from the vector of Fig. 1 by treatment of the vector with restriction endonucleases appropriate for removal of the supplemental sequences. The supplemental sequence subsequent to the portable DNA sequence, identified in Fig. 1 as region C, may be removed by treatment of the vector with a suitable restriction endonuclease, preferably *HgiA1* followed by reconstruction of the 3' end of region A using synthetic oligonucleotides and ligation of the vector with T4 DNA ligase. Deletion of the supplemental sequence preceding the portable DNA sequence, identified as region B in Fig. 1, would be specifically accomplished by the method set forth in Example 2.

[045] In preferred embodiments, cloning vectors containing and capable of expressing the portable DNA sequence of the present invention contain various operational elements. These "operational elements," as discussed herein, include at least one promoter, at least one Shine-Dalgarno sequence, at least one terminator codon. Preferably, these "operational elements" also include at least one operator, at least one leader sequence, and for proteins to be exported from intracellular space, at least one regulator and any other DNA sequences necessary or preferred for appropriate transcription and subsequent translation of the vector DNA.

[046] Additional embodiments of the present invention are envisioned as employing other known or currently undiscovered vectors which would contain one or

more of the portable DNA sequences described herein. In particular, it is preferred that these vectors have some or all of the following characteristics: (1) possess a minimal number of host-organism sequences; (2) be stable in the desired host; (3) be capable of being present in a high copy number in the desired host; (4) possess a regulatable promoter; (5) have at least one DNA sequence coding for a selectable trait present on a portion of the plasmid separate from that where the portable DNA sequence will be inserted; and (6) be integrated into the vector.

[047] The following, noninclusive, list of cloning vectors is believed to set forth vectors which can easily be altered to meet the above criteria and are therefore preferred for use in the present invention. Such alterations are easily performed by those of ordinary skill in the art in light of the available literature and the teachings herein.

TABLE I

HOST	Vectors	Comments
<u>E. coli</u>	pUC8 pUC9 pBR322 pGW7 placI ^q pDP8	Many selectable replicons have been characterized. Maniatis, T. <i>et al.</i> (1982), <u>Molecular Cloning: A Laboratory Manual</u> , Cold Spring Harbor Laboratory.
<u>BACILLUS</u> <u>B. subtilis</u> <u>B. amyloliquefaciens</u> <u>B. stearothermophilus</u>	pUB110 pSA0501 pSA2100 pBD6 pBD8 pT127	<u>Genetics and Biotechnology of Bacilli</u> , Ganesan and Hoch, eds., 1984, Academic Press.
<u>PSEUDOMONAS</u> <u>P. aeruginosa</u> <u>P. putida</u>	RSF1010 Rms149 pKT209 RK2 pSa727	Some vectors useful in broad host range of gram - negative bacteria including Xanthomonas and Agrobacterium

CLOSTRIDIUM

C. perfringens

pJU12

pJU7

pJU10

pJU16

pJU13

SACCHAROMYCES

S. cerevisiae

YEp24

YIp5

YRp17

Shuttle plasmids for *E. coli* and *C. perfringens* construction ref. Squires, C. et al. (1984) Journal Bacteriol. 159:465-471. Botstein and Davis in Molecular Biology of the Yeast Saccharomyces, Strathern, Jones, and Broach, eds., 1982, Cold Spring Harbor Laboratory.

[048] It is to be understood that additional cloning vectors may now exist or will be discovered which have the above-identified properties and are therefore suitable for use in the present invention. These vectors are also contemplated as being within the scope of the disclosed series of cloning vectors into which the portable DNA sequences may be introduced, along with any necessary operational elements, and which altered vector is then included within the scope of the present invention and would be capable of being used in the recombinant-DNA method set forth more fully below.

[049] In addition to the above list, an *E. coli* vector system, as set forth in Example 2, is preferred in one embodiment as a cloning vector. Moreover, several vector plasmids which autonomously replicate in a broad range of Gram Negative bacteria are preferred for use as cloning vehicles in hosts of the genera *Pseudomonas*. These are described by Tait, R.C., Close, T.J., Lundquist, R.C., Hagiya, M., Rodriguez, R.L., and Kado, C.I. in *Biotechnology*, May, 1983, pp. 269-275; Panopoulos, N.J. in Genetic Engineering in the Plant Sciences, Praeger Publishers, New York, New York, pp. 163-185, (1981); and Sakaguchi, K. in *Current Topic in Microbiology and Immunology* 96:31-45, (1982), each of which is specifically incorporated herein by reference.

[050] One particularly preferred construction employs the plasmid RSF1010 and derivatives thereof as described by Bagdasarian, M., Bagdasarian, M.M., Coleman, S., and Timmis, K.N. in Plasmids of Medical Environmental and Commercial Importance, Timmis, K.N. and Puhler, A. eds., Elsevier/North Holland Biomedical Press, (1979), specifically incorporated herein by reference. The advantages of RSF1010 are that it is relatively small, high copy number plasmid which is readily transformed into and stably maintained in both *E. coli* and *Pseudomonas* species. In this system, it is preferred to use the Tac expression system as described for *Escherichia*, since it appears that the *E. coli* trp promoter is readily recognized by *Pseudomonas* RNA polymerase as set forth by Sakaguchi, K. in Current Topics in Microbiology and Immunology 96:31-45 (1982) and Gray, G.L., McKeown, K.A., Jones, A.J.S., Seeburg, P.H., and Heyneker, H.L. in Biotechnology Feb. 1984, pp. 161-165, both of which are specifically incorporated herein by reference. Transcriptional activity may be further maximized by requiring the exchange of the promoter with, e.g., an *E. coli* or *P. aeruginosa* trp promoter.

[051] In a preferred embodiment, *P. aeruginosa* is transformed with vectors directing the synthesis of the metalloproteinase inhibitor as either an intracellular product or as a product coupled to leader sequences that will effect its processing and export from the cell. In this embodiment, these leader sequences are preferably selected from the group consisting of beta-lactamase, OmpA protein, the naturally occurring human signal peptide, and that of carboxypeptidase G2 from *Pseudomonas*. Translation may be coupled to translation initiation for any of the *E. coli* proteins as

described in Example 2, as well as to initiation sites for any of the highly expressed proteins of the host to cause intracellular expression of the metalloproteinase inhibitor.

[052] In those cases where restriction minus strains of a host *Pseudomonas* species are not available, transformation efficiency with plasmid constructs isolated from *E. coli* are poor. Therefore, passage of the *Pseudomonas* cloning vector through an r- m+ strain of another species prior to transformation of the desired host, as set forth in Bagdasarian, M., *et al.*, Plasmids of Medical, Environmental and Commercial Importance, pp. 411-422, Timmis and Puhler eds., Elsevier/North Holland Biomedical Press (1979), specifically incorporated herein by reference, is desired.

[053] Furthermore, a preferred expression system in hosts of the genera *Bacillus* involves using plasmid pUB110 as the cloning vehicle. As in other host vector systems, it is possible in *Bacillus* to express the metalloproteinase inhibitors of the present invention as either an intracellular or a secreted protein. The present embodiments include both systems. Shuttle vectors that replicate in both *Bacillus* and *E. coli* are available for constructing and testing various genes as described by Dubnau, D., Gryczan, T., Contente, S., and Shivakumar, A.G. in Genetic Engineering, Vol. 2, Setlow and Hollander eds., Plenum Press, New York, New York, pp. 115-131, (1980), specifically incorporated herein by reference. For the expression and secretion of metalloproteinase inhibitors from *B. subtilis*, the signal sequence of alpha-amylase is preferably coupled to the coding region for the metalloproteinase inhibitor. For synthesis of intracellular metalloproteinase inhibitor, the portable DNA sequence will be translationally coupled to the ribosome binding site of the alpha-amylase leader sequence.

[054] Transcription of either of these constructs is preferably directed by the alpha-amylase promoter or a derivative thereof. This derivative contains the RNA polymerase recognition sequence of the native alpha-amylase promoter but incorporates the lac operator region as well. Similar hybrid promoters constructed from the penicillinase gene promoter and the lac operator have been shown to function in *Bacillus* hosts in a regulatable fashion as set forth by Yansura, D.G. and Henner in Genetics and Biotechnology of Bacilli, Ganesan, A.T. and Hoch, J.A., eds., Academic Press, pp. 249-263, (1984), specifically incorporated by reference. The *lacI*^q would also be included to effect regulation.

[055] One preferred construction for expression in *Clostridium* is in plasmid pJU12 described by Squires, C. H. *et al.* in J. Bacteriol. 159:465-471 (1984), specifically incorporated herein by reference, transformed into *C. perfringens* by the method of Heefner, D. L. *et al.* as described in J. Bacteriol. 159:460-464 (1984), specifically incorporated herein by reference.

[056] Transcription is directed by the promoter of the tetracycline resistance gene. Translation is coupled to the Shine-Dalgarno sequences of this same *tet*^r gene in a manner strictly analogous to the procedures outlined above for vectors suitable for use in other hosts.

[057] Maintenance of foreign DNA introduced into yeast can be effected in several ways (Botstein, D., and Davis, R. W., in The Molecular Biology of the Yeast *Saccharomyces*, Cold Spring Harbor Laboratory, Strathern, Jones and Broach, eds., pp. 607-636 (1982)). One preferred expression system for use with host organisms of the genus *Saccharomyces* harbors the anticollagenase gene on the 2 micron plasmid.

The advantages of the 2 micron circle include relatively high copy number and stability when introduced into *cir^o* strains. These vectors preferably incorporate the replication origin and at least one antibiotic resistance marker from pBR322 to allow replication and selection in *E. coli*. In addition, the plasmid will preferably have 2 micron sequences and the yeast LEU2 gene to serve the same purposes in LEU2 mutants of yeast.

[058] The regulatable promoter from the yeast *GALI* gene will preferably be adapted to direct transcription of the portable DNA sequence gene. Translation of the portable DNA sequence in yeast will be coupled to the leader sequence that directs the secretion of yeast alpha-factor. This will cause formation of a fusion protein which will be processed in yeast and result in secretion of a metalloproteinase inhibitor. Alternatively, a methionylmetalloproteinase inhibitor will be translated for inclusion within the cell.

[059] It is anticipated that translation of mRNA coding for the metalloproteinase inhibitor in yeast will be more efficient with the preferred codon usage of yeast than with the sequence present in pUC8-Fic, as identified in Example 2, which has been tailored to the prokaryotic bias. For this reason, the portion of the 5' end of the portable DNA sequence beginning at the *Tth111I* site is preferably resynthesized.

505370-454860
SUBC8) The new sequence favors the codons most frequently used in yeast. This new sequence preferably has the following nucleotide sequence:

HgiAI

```

5'   GAT  CCG  TGC  ACT  TGT  GTT  CCA  CCA  CAC
      GC  ACG  TGA  ACA  CAA  GGT  GGT  GTG

      CCA  CAA  ACT  GCT  TTC  TGT  AAC  TCT  GAC  C
      GGT  GTT  TGA  CGA  AAG  ACA  TTG  AGA  CTG  GA  3'
  
```

[060] As will be seen from an examination of the individual cloning vectors and systems contained on the above list and description, various operational elements may be present in each of the preferred vectors of the present invention. It is contemplated any additional operational elements which may be required may be added to these vectors using methods known to those of ordinary skill in the art, particularly in light of the teachings herein.

[061] In practice, it is possible to construct each of these vectors in a way that allows them to be easily isolated, assembled, and interchanged. This facilitates assembly of numerous functional genes from combinations of these elements and the coding region of the metalloproteinase inhibitor. Further, many of these elements will be applicable in more than one host.

[062] At least one origin of replication recognized by the contemplated host microorganism, along with at least one selectable marker and at least one promoter sequence capable of initiating transcription of the portable DNA sequence are contemplated as being included in these vectors. It is additionally contemplated that the vectors, in certain preferred embodiments, will contain DNA sequences capable of functioning as regulators ("operators"), and other DNA sequences capable of coding for regulator proteins. In preferred vectors of this series, the vectors additionally contain ribosome binding sites, transcription terminators and leader sequences.

[063] These regulators, in one embodiment, will serve to prevent expression of the portable DNA sequence in the presence of certain environmental conditions and, in the presence of other environmental conditions, allow transcription and subsequent expression of the protein coded for by the portable DNA sequence. In particular, it is

preferred that regulatory segments be inserted into the vector such that expression of the portable DNA sequence will not occur in the absence of, for example, isopropylthio- β -d-galactoside. In this situation, the transformed microorganisms containing the portable DNA may be grown to a desired density prior to initiation of the expression of the metalloproteinase inhibitor. In this embodiment, expression of the desired protease inhibitor is induced by addition of a substance to the microbial environment capable of causing expression of the DNA sequence after the desired density has been achieved.

[064] Additionally, it is preferred that an appropriate secretory leader sequence be present, either in the vector or at the 5' end of the portable DNA sequence, the leader sequence being in a position which allows the leader sequence to be immediately adjacent to the initial portion of the nucleotide sequence capable of directing expression of the protease inhibitor without any intervening transcription or translation termination signals. The presence of the leader sequence is desired in part for one or more of the following reasons: 1) the presence of the leader sequence may facilitate host processing of the initial product to the mature recombinant metalloproteinase inhibitor; 2) the presence of the leader sequence may facilitate purification of the recombinant metalloproteinase inhibitors, through directing the metalloproteinase inhibitor out of the cell cytoplasm; 3) the presence of the leader sequence may affect the ability of the recombinant metalloproteinase inhibitor to fold to its active structure through directing the metalloproteinase inhibitor out of the cell cytoplasm.

[065] In particular, the leader sequence may direct cleavage of the initial translation product by a leader peptidase to remove the leader sequence and leave a polypeptide with the amino acid sequence which has the potential of metalloproteinase inhibitory activity. In some species of host microorganisms, the presence of the appropriate leader sequence will allow transport of the completed protein into the periplasmic space, as in the case of *E. coli*. In the case of certain yeasts and strains of *Bacillus* and *Pseudomonas*, the appropriate leader sequence will allow transport of the protein through the cell membrane and into the extracellular medium. In this situation, the protein may be purified from extracellular protein.

[066] Thirdly, in the case of some of the metalloproteinase inhibitors prepared by the present invention, the presence of the leader sequence may be necessary to locate the completed protein in an environment where it may fold to assume its active structure, which structure possesses the appropriate metalloproteinase activity.

[067] Additional operational elements include, but are not limited to, ribosome-binding sites and other DNA sequences necessary for microbial expression of foreign proteins. The operational elements as discussed herein can be routinely selected by those of ordinary skill in the art in light of prior literature and the teachings contained herein. General examples of these operational elements are set forth in B. Lewin, Genes, Wiley & Sons, New York (1983), which is specifically incorporated herein by reference. Various examples of suitable operational elements may be found on the vectors discussed above and may be elucidated through review of the publications discussing the basic characteristics of the aforementioned vectors.

[068] In one preferred embodiment of the present invention, an additional DNA sequence is located immediately preceding the portable DNA sequence which codes for the metalloproteinase inhibitor. The additional DNA sequence is capable of functioning as a translational coupler, *i.e.*, it is a DNA sequence that encodes an RNA which serves to position ribosomes immediately adjacent to the ribosome binding site of the metalloproteinase inhibitor RNA with which it is contiguous.

[069] Upon synthesis and/or isolation of all necessary and desired component parts of the above-discussed cloning vectors, the vectors are assembled by methods generally known to those of ordinary skill in the art. Assembly of such vectors is believed to be within the duties and tasks performed by those with ordinary skill in the art and, as such, is capable of being performed without undue experimentation. For example, similar DNA sequences have been ligated into appropriate cloning vectors, as set forth in Schoner *et al.*, Proceedings of the National Academy of Sciences U.S.A., 81:5403-5407 (1984), which is specifically incorporated herein by reference.

[070] In construction of the cloning vectors of the present invention, it should additionally be noted that multiple copies of the portable DNA sequence and its attendant operational elements may be inserted into each vector. In such an embodiment, the host organism would produce greater amounts per vector of the desired metalloproteinase inhibitor. The number of multiple copies of the DNA sequence which may be inserted into the vector is limited only by the ability of the resultant vector, due to its size, to be transferred into and replicated and transcribed in an appropriate host microorganism.

[071] Additionally, it is preferred that the cloning vector contain a selectable marker, such as a drug resistance marker or other marker which causes expression of a selectable trait by the host microorganism. In a particularly preferred embodiment of the present invention, the gene for ampicillin resistance is included in vector pUC9-F5/237P10.

[072] Such a drug resistance or other selectable marker is intended in part to facilitate in the selection of transformants. Additionally, the presence of such a selectable marker on the cloning vector may be of use in keeping contaminating microorganisms from multiplying in the culture medium. In this embodiment, such a pure culture of the transformed host microorganisms would be obtained by culturing the microorganisms under conditions which require the induced phenotype for survival.

[073] It is noted that, in preferred embodiment, it is also desirable to reconstruct the 3' end of the coding region to allow assembly with 3' non-translated sequences. Included among these non-translated sequences are those which stabilize the mRNA or enhance its transcription and those that provide strong transcriptional termination signals which may stabilize the vector as they are identified by Gentz, R., Langner, A., Chang, A.C.Y., Cohen, S.H., and Bujard, H. in Proc. Natl. Acad. Sci. USA 78: 4936-4940 (1981), specifically incorporated herein by reference.

[074] This invention also relates to a recombinant-DNA method for the production of metalloproteinase inhibitors. Generally, this method includes:

- (a) preparation of a portable DNA sequence capable of directing a host microorganism to produce a protein having metalloproteinase inhibitor activity;

- (b) cloning the portable DNA sequence into a vector capable of being transferred into and replicating in a host microorganism, such vector containing operational elements for the portable DNA sequence;
- (c) transferring the vector containing the portable DNA sequence and operational elements into a host microorganism capable of expressing the metalloproteinase inhibitor protein;
- (d) culturing the host microorganism under conditions appropriate for amplification of the vector and expression of the inhibitor; and
- (e) in either order:
- (i) harvesting the inhibitor; and
 - (ii) causing the inhibitor to assume an active, tertiary structure whereby it possesses metalloproteinase inhibitor activity.

[075] In this method, the portable DNA sequences are those synthetic or naturally-occurring polynucleotides described above. In a preferred embodiment of the present method, the portable DNA sequence has the nucleotide sequence as follows:

10 20 30 40 50 60
GTTGTTGCTG TGGCTGATAG CCCCAGCAGG GCCTGCACCT GTGTCCCACC CCACCCACAG

70 80 90 100 110 120
ACGGCCTTCT GCAATTCCGA CCTCGTCATC AGGGCCAAGT TCGTGGGGAC ACCAGAAGTC

130 140 150 160 170 180
AACCAGACCA CTTTATACCA GCGTTATGAG ATCAAGATGA CCAAGATGTA TAAAGGGTTC

190 200 210 220 230 240
CAAGCCTTAG GGGATGCCGC TGACATCCGG TTCGTCTACA CCCCCGCCAT GGAGAGTGTC

250 260 270 280 290 300
TGCGGATACT TCCACAGGTC CCACAACCGC AGCGAGGAGT TTCTCATTGC TGGAAAAGTC

310 320 330 340 350 360
CAGGATGGAC TCTTGACAT CACTACCTGC AGTTTCGTGG CTCCCTGGAA CAGCCTGAGC

370 380 390 400 410 420
 TTAGCTCAGC GCCGGGGCTT CACCAAGACC TACTGTGTTG GCTGTGAGGA ATGCACAGTG
 430 440 450 460 470 480
 TTTCCCTGTT TATCCATCCC CTGCAAACTG CAGAGTGGCA CTCATTGCTT GTGGACGGAC
 490 500 510 520 530 540
 CAGCTCCTCC AAGGCTCTGA AAAGGGCTTC CAGTCCCGTC ACCTTGCCTG CCTGCCTCGG
 550 560 570 580 590 600
 GAGCCAGGGC TGTGCACCTG GCAGTCCCTG CGGTCCCAGA TAGCCTGAAT CCTGCCCGGA
 610 620 630 640 650 660
 GTGGAAGCTG AAGCCTGCAC AGTGTCCACC CTGTTCCAC TCCCATCTTT CTTCCGGACA
 670 680 690 700
 ATGAAATAAA GAGTTACCAC CCAGCAAAAA AAAAAAGGAA TTC

[076] The vectors contemplated as being useful in the present method are those described above. In a preferred embodiment, the cloning vector pUC9-F5/237P10 is used in the disclosed method.

[077] The vector thus obtained is then transferred into the appropriate host microorganism. It is believed that any microorganism having the ability to take up exogenous DNA and express those genes and attendant operational elements may be chosen. It is preferred that the host microorganism be an anaerobe, facultative anaerobe or aerobe. Particular hosts which may be preferable for use in this method include yeasts and bacteria. Specific yeasts include those of the genus *Saccharomyces*, and especially *Saccharomyces cerevisiae*.

[078] Specific bacteria include those of the genera *Bacillus* and *Escherichia* and *Pseudomonas*. Various other preferred hosts are set forth in Table I, *supra*. In other, alternatively preferred embodiments of the present invention, *Bacillus subtilis*, *Escherichia coli* or *Pseudomonas aeruginosa* are selected as the host microorganisms.

[079] After a host organism has been chosen, the vector is transferred into the host organism using methods generally known by those of ordinary skill in the art. Examples of such methods may be found in Advanced Bacterial Genetics by R. W. Davis *et al.*, Cold Spring Harbor Press, Cold Spring Harbor, New York, (1980), which is specifically incorporated herein by reference. It is preferred, in one embodiment, that the transformation occur at low temperatures, as temperature regulation is contemplated as a means of regulating gene expression through the use of operational elements as set forth above. In another embodiment, if osmolar regulators have been inserted into the vector, regulation of the salt concentrations during the transformation would be required to insure appropriate control of the synthetic genes.

[080] If it is contemplated that the recombinant metalloproteinase inhibitors will ultimately be expressed in yeast, it is preferred that the cloning vector first be transferred into *Escherichia coli*, where the vector would be allowed to replicate and from which the vector would be obtained and purified after amplification. The vector would then be transferred into the yeast for ultimate expression of the metalloproteinase inhibitor.

[081] The host microorganisms are cultured under conditions appropriate for the expression of the metalloproteinase inhibitor. These conditions are generally specific for the host organism, and are readily determined by one of ordinary skill in the art, in light of the published literature regarding the growth conditions for such organisms, for example Bergey's Manual of Determinative Bacteriology, 8th Ed., Williams & Wilkins Company, Baltimore, Maryland, which is specifically incorporated herein by reference.

09987357 "043602 205270 2572860

[082] Any conditions necessary for the regulation of the expression of the DNA sequence, dependent upon any operational elements inserted into or present in the vector, would be in effect at the transformation and culturing stages. In one embodiment, the cells are grown to a high density in the presence of appropriate regulatory conditions which inhibit the expression of the DNA sequence. When optimal cell density is approached, the environmental conditions are altered to those appropriate for expression of the portable DNA sequence. It is thus contemplated that the production of the metalloproteinase inhibitor will occur in a time span subsequent to the growth of the host cells to near optimal density, and that the resultant metalloproteinase inhibitor will be harvested at some time after the regulatory conditions necessary for its expression were induced.

[083] In a preferred embodiment of the present invention, the recombinant metalloproteinase inhibitor is purified subsequent to harvesting and prior to assumption of its active structure. This embodiment is preferred as the inventors believe that recovery of a high yield of re-folded protein is facilitated if the protein is first purified. However, in one preferred, alternate embodiment, the metalloproteinase inhibitor may be allowed re-fold to assume its active structure prior to purification. In yet another preferred, alternate embodiment, the metalloproteinase inhibitor is caused to assume its re-folded, active state upon recovery from the culturing medium.

SUB C10) [084] In certain circumstances, the metalloproteinase inhibitor will assume its proper, active structure upon expression in the host microorganism and transport of the protein through the cell wall or membrane or into the periplasmic space. This will generally occur if DNA coding for an appropriate leader sequence has been linked to

the DNA coding for the recombinant protein. The preferred metalloproteinase inhibitors of the present invention will assume their mature, active form upon translocation out of the inner cell membrane. The structures of numerous signal peptides have been published, for example by Marion E.E. Watson in Nuc. Acid Res. 12: 515-5164, 1984, specifically incorporated herein by reference. It is intended that these leader sequences, together with portable DNA, will direct intracellular production of a fusion protein which will be transported through the cell membrane and will have the leader sequence portion cleaved upon release from the cell.

[085] In a preferred embodiment, the signal peptide of the *E. coli* OmpA protein is used as a leader sequence and is located in a position contiguous with the portable DNA sequence coding for the metalloproteinase inhibitor structure.

[086] Additionally preferred leader sequences include those of beta-lactamase, carboxypeptidase G2 and the human signal protein. These and other leader sequences are described.

[087] If the metalloproteinase inhibitor does not assume its proper, active structure, any disulfide bonds which have formed and/or any noncovalent interactions which have occurred will first be disrupted by denaturing and reducing agents, for example, guanidinium chloride and β -mercaptoethanol, before the metalloproteinase inhibitor is allowed to assume its active structure following dilution and oxidation of these agents under controlled conditions.

[088] The transcription terminators contemplated herein serve to stabilize the vector. In particular, those sequences as described by Gentz *et al.*, in Proc. Natl. Acad.

Sub C10
Cont

205270-4528650

Sci. USA 78: 4936-4940 (1981), specifically incorporated herein by reference, are contemplated for use in the present invention.

[089] It is to be understood that application of the teachings of the present invention to a specific problem or environment will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein. Examples of the products of the present invention and representative processes for their isolation and manufacture appear in the following examples.

EXAMPLES

EXAMPLE 1

Preparation of Poly(A⁺) RNA from HEF-SA Fibroblasts

[090] HEF-SA cells were grown to near confluence in 75 cm² T-flasks. Cells were washed twice in Dulbecco's phosphate buffered saline solution and harvested by the addition of 2 ml of 10 mM Tris, pH 7.5 containing 1% w/v SDS (obtained from BDH Chemicals, Ltd., Poole, England), 5 mM EDTA and 20 µg/ml protease K (obtained from Boehringer Mannheim Biochemicals, Indianapolis, Indiana). Each flask was subsequently washed with an additional milliliter of this same solution.

[091] The pooled aliquots from the cell harvest were made to 70 µg/ml in protease K and incubated at 40°C for 45 minutes. The proteolyzed solution was brought to a NaCl concentration of 150 mM by the addition of 5 M stock and subsequently extracted with an equal volume of phenol:chloroform 1:1. The aqueous phase was reextracted with an equal volume of chloroform. Two volumes of ethanol were added to the aqueous phase and incubated overnight at -20°C. The precipitated nucleic acids were recovered by centrifugation at 17,500 xg for 10 minutes in a Beckman J2-21 centrifuge, Beckman Instruments, Palo Alto, California, and were

redissolved in 25 ml of 0.1% w/v SDS. This solution was again extracted with an equal volume of chloroform. The aqueous phase was added to two volumes of cold ethanol and kept at -20°C for 2 hours. The precipitate was collected by centrifugation at 10,000 xg for 15 minutes and redissolved in 10 ml of 1 mM Tris, 0.5 mM EDTA, 0.1% SDS, pH 7.5. RNA was precipitated from this solution by the addition of 10 ml of 4 M LiCl, 20 mM NaOAc, pH 5.0 and incubated at -20°C for 18 hours. The precipitate was again recovered by centrifugation and washed twice with 2 M LiCl before redissolving in 1 mM Tris, 0.5 mM EDTA, 0.1% SDS, pH 7.5. This solution was stored at -70°C.

Chromatography on Oligo dT Cellulose

[092] Total cellular RNA prepared as above was ethanol precipitated and redissolved in 0.5 M NaCl. Five ml of RNA at 0.45 mg/ml were applied to a 1 ml column of washed type VII oligo dT cellulose (obtained from PL Biochemicals, Milwaukee, Wisconsin). The column was then washed with 10 ml of 0.5 M NaCl and eluted with 2.0 ml of sterile H₂O. The eluted poly(A⁺) fraction of RNA was ethanol precipitated and dissolved to give a 1 mg/ml solution in 1 mM Tris, 0.1 mM EDTA, pH 8.0. This was stored at -70°C.

cDNA Synthesis

[093] Poly(A⁺) RNA was primed with oligo dT (obtained from PL Biochemicals, Milwaukee, Wisconsin) to serve as a template for cDNA synthesis by AMV reverse transcriptase (obtained from Life Sciences, Inc., St. Petersburg, Florida). Following the synthesis reaction, the RNA was hydrolyzed by the addition of 0.1 volume of 3 N NaOH and incubated at 67°C for 10 minutes. The solution was then neutralized and the cDNA purified by gel filtration chromatography on biogel A 1.5 (obtained from BioRad

Laboratories, Richmond, California) in a 0.7x25 cm column in a 10 mM Tris, 5 mM EDTA, and 1% SDS, pH 7.5 solution. Fractions containing cDNA were pooled and concentrated by ethanol precipitation. The cDNA was dG tailed and purified by gel filtration using the procedure set forth above. Second strand synthesis was primed with oligo dC and polymerized in an initial reaction with the large (Klenow) fragment of DNA polymerase (obtained from Boehringer Mannheim). Following second strand synthesis, *E. coli* DNA polymerase I (obtained from Boehringer Mannheim) was added and incubation continued to form blunt ends. The double stranded cDNA was again purified by chromatography. *Eco*RI restriction sites within the cDNAs were modified by the action of *Eco*RI methylase, obtained from New England Biolabs, Beverly, Massachusetts. The cDNA was again purified and ligated to synthetic *Eco*RI linkers. Finally, the ends were then trimmed with the endonuclease and the cDNA purified by gel filtration. This DNA was ligated into a unique *Eco*RI site in lambda gt10 DNA packaged *in vitro* and used to infect *E. coli* strain hflA according to the method set forth by Huynh, T.V., Young, R.A., and Davis, R.W., in DNA Cloning Techniques, A Practical Approach (ed. Glover, D.M.) (IRL Press Oxford), in press, specifically incorporated herein by reference. Approximately 25,000 recombinants were amplified in this manner.

Screening

[094] Recombinant-phage-containing sequences of interest were selected by their preferential hybridization to synthetic oligonucleotides encoding portions of the primary structure of the desired metalloproteinase inhibitor, hereinafter referred to as FIBAC. These portions of the protein sequence correspond in part to those set forth in

the published literature by Stricklin, G.P. and Welgus, H.G., J. Biol. Chem. 258: 12252-12258 (1983), specifically incorporated herein by reference. Recombinant phage were used to infect *E. coli* strain hflA and plated at a density of approximately 2×10^3 pfu/150 mm petri dish. Phage were blotted onto nitrocellulose filters (BA85, Schleicher & Schuell Inc., Keene, New Hampshire), and DNA was denatured and fixed essentially as described by Benton and Davis in Science 196: 180-182 (1979) specifically incorporated herein by reference.

[095] Using that procedure, the filters were treated sequentially for 10-15 minutes each in 0.5 M NaCl, then 1.0 M Tris, 1.5 M NaCl pH 8.0, and finally submerged in 2x SSPE. (2x SSPE is 0.36 M NaCl, 20 mM NaH_2PO_4 , 2 mM EDTA, pH 7.4). Filters were blotted dry and baked 75°-80° for 3-4 hours. Duplicate filters were made of each plate. Filters were prehybridized for 1-3 hours at 37° in 5x SSPE containing 0.1x SET, 0.15% NaPPi, and 1x Denhardts solutions. Filters were then hybridized for 72 hours at 37° in this same solution containing 5×10^5 cpm/ml of 5' end-labeled 51-mer oligonucleotide (specific activity approximately 10^6 cpm/pmole). Following hybridization, filters were washed six times in 5x SSPE containing 0.1x SET and 0.05% sodium pyrophosphate at 37°, then three times in 2x SSPE at 21°. These were then blotted dry and autoradiographed on Kodak XAR-5 film at -70° with a Kodak lightening-plus intensifying screen. Signals clearly visible from duplicate filters were used to pick phage for plaque purification. Filter preparations and hybridization procedures for plaque purification steps were the same as above. The washing procedure was simplified to 6 changes of 2x SSPE at 37°. Six isolates purified by repetitive plating

lambda FIBAC 5 on agarose gel electrophoresis. This plasmid has been named pUC9-F5/237P10.

Mapping and Subcloning

[098] The insert in pUC9-F5/237P10 was mapped with respect to internal *Pst*I sites. Double digests with *Eco*RI and *Pst*I demonstrated three internal *Pst*I recognition sites. The entire insert and the component pieces were subcloned into M13 bacteriophage mp19 and mp18, respectively. Sequencing of the pieces was performed by the dideoxynucleotide method described by Sanger *et al.* in Sanger, F., Nicklen, S., and Coulson, A.R., Proc. Natl. Acad. Sci. USA 74: 5463-5467 (1977), specifically incorporated herein by reference.

[099] The sequence of the DNA insert from pUC9-F5/237P10 showed an open reading frame which encodes the primary structure of a mature fibroblast collagenase inhibitor biologically equivalent to that isolable from human skin fibroblasts. The salient features of the sequence are:

- (1) The insert is flanked by *Eco*RI restriction sites and by G/C and A/T homopolymeric tracts consistent with the cloning methodology;
- (2) The coding strand is presented in the 5' to 3' convention with poly C at the 5' end and poly A at the 3' end, again consistent with the techniques employed;
- (3) If the first G in the sequence GTTGTTG immediately adjacent to the 3' end of the poly C tract is considered as nucleotide 1, then an open reading frame is presented which encodes the primary structure of the

mature human fibroblast collagenase inhibitor beginning at nucleotide 34 and continuing through nucleotide 585;

- (4) The termination codon TGA at nucleotides 586 through 588 defines the carboxy terminus of the translation product which is the same as that of the mature protein;
- (5) Nucleotides 1 through 33 define an amino acid sequence which is not found in the primary structure of the processed protein, but which is probably a portion of a leader peptide characteristic of secreted proteins;
- (6) The three internal *Pst*I sites have as their first base nucleotides 298, 327, and 448;
- (7) There is a single recognition sequence for the restriction enzyme *Tth*111I beginning at nucleotide 78; and
- (8) There is a single recognition sequence for the restriction endonuclease *Nco*I beginning at nucleotide 227.

[0100] The sequence of nucleotides 1 through 703 and restriction site analysis are shown.

	#	SITES	FRAGMENTS		FRAGMENTS	ENDS
ACC 1 (GTVWAC)	1	214	495	(69.8)	214	709
			214	(30.2)	1	214
ALU 1 (AGCT)	4	358	358	(50.5)	1	358
		363	124	(17.5)	482	606
		482	119	(16.8)	363	482
		606	103	(14.5)	606	709
			5	(0.7)	358	363
AVA 1 (CQCGPG)	1	536	536	(75.6)	1	536
			173	(24.4)	536	709

	#	SITES	FRAGMENTS	FRAGMENTS	ENDS
AVA 2 (GGRCC)	3	257	257 (36.2)	1	257
		477	220 (31.0)	257	477
		572	137 (19.3)	572	709
			95 (13.4)	477	572
BBV 1 (GCTGC)	1	269	440 (62.1)	269	709
			269 (37.9)	1	269
BST N1 (CCRGG)	3	344	344 (48.5)	1	344
		544	200 (28.2)	344	544
		557	152 (21.4)	557	709
			13 (1.8)	544	557
DDE 1 (CTNAG)	4	186	344 (48.5)	365	709
		355	186 (26.2)	1	186
		360	169 (23.8)	186	355
	#	SITES	FRAGMENTS	FRAGMENTS	ENDS
		365	5 (0.7)	360	365
			5 (0.7)	355	360
ECO R1 (GAATTC)	1	698	698 (98.4)	1	698
			11 (1.6)	698	709
FNU4H 1 (GCNGC)	2	196	440 (62.1)	269	709
		269	196 (27.6)	1	196
			73 (10.3)	196	269
FOK 1 (GGATG)	4	192	274 (38.6)	435	709
		204	192 (27.1)	1	192
		303	132 (18.6)	303	435
		435	99 (14.0)	204	303
			12 (1.7)	192	204
HAE 2 (PGCGCQ)	1	368	368 (51.9)	1	368
			341 (48.1)	368	709

	#	SITES	FRAGMENTS	FRAGMENTS	ENDS
HAE 3 (GGCC)	3	30 63 93	616 (86.9) 33 (4.7) 30 (4.2) 30 (4.2)	93 30 63 1	709 63 93 30
HGI A1 (GRGCRC)	1	552	552 (77.9) 157 (22.1)	1 552	552 709
HHA 1 (GCGC)	1	369	369 (52.0) 340 (48.0)	1 369	369 709
HINC 2 (GTQPAC)	1	118	591 (83.4) 118 (16.6)	118 1	709 118
HINF 1 (GANTC)	2	308 587	308 (43.4) 279 (39.4) 122 (17.2)	1 308 587	308 587 709
HPA 2 (CCGG)	4	207 372 596 654	224 (31.6) 207 (29.2) 165 (23.3) 58 (8.2) 55 (7.8)	372 1 207 596 654	596 207 372 654 709
HPH 1 (GGTGA)	2	380 519	380 (53.6) 190 (26.8) 139 (19.6)	1 519 380	380 709 519
MBO 2 (GAAGA)	1	650	650 (91.7) 59 (8.3)	1 650	650 709

	#	SITES	FRAGMENTS		FRAGMENTS	ENDS
MNL 1 (CCTC)	5	81	193	(27.2)	81	274
		274	174	(24.5)	535	709
		406	132	(18.6)	274	406
		486	81	(11.4)	1	81
		535	80	(11.3)	406	486
			49	(6.9)	486	535
MST 2 (CCTNAGG)	1	185	524	(73.9)	185	709
			185	(26.1)	1	185
NCI 1 (CCSGG)	2	372	372	(52.5)	1	372
		595	223	(31.5)	372	595
			114	(16.1)	595	709
NCO 1 (CCATGG)	1	227	482	(68.0)	227	709
			227	(32.0)	1	227
NSP B2 (CVGCWG)	1	197	512	(72.2)	197	709
			197	(27.8)	1	197
PST 1 (CTGCAG)	3	298	298	(42.0)	1	298
		327	261	(36.8)	448	709
		448	121	(17.1)	327	448
			29	(4.1)	298	327
SAU 1 (CCTNAGG)	1	185	524	(73.9)	185	709
			185	(26.1)	1	185
SAU 3A (GATC)	1	150	559	(78.8)	150	709
			150	(21.2)	1	150
SAU96 1 (GGNCC)	5	29	220	(31.0)	257	477
		92	165	(23.3)	92	257
		257	137	(19.3)	572	709
		477	95	(13.4)	477	572
		572	63	(8.9)	29	92
			29	(4.1)	1	29

	#	SITES	FRAGMENTS	FRAGMENTS	ENDS
SCR F1 (CCNGG)	5				
		344	344 (48.5)	1	344
		372	172 (24.3)	372	544
		544	114 (16.1)	595	709
		557	38 (5.4)	557	595
		595	28 (3.9)	344	372
			13 (1.8)	544	557

SFA N1 (GATGC)	1				
		193	516 (72.8)	193	709
			193 (27.2)	1	193

TTH111 I	1				
(GACNNNGTC)		79	630 (88.9)	79	709
			79 (11.1)	1	79

[0101] The following do not appear:

AAT 2	AFL 2	AFL 3	AHA 3
APA 1	ASU 2	AVA 3	AVR 2
BAL 1	BAM H1	BCL 1	BGL 1
BGL 2	BIN 1	BSSH 1	BST E2
CFR 1	CLA 1	ECO R5	FNUD 2
GDI 2	HAE 1	HGA 1	HGI C1
HGI D1	HGI J2	HIND 3	HPA 1
KPN 1	MLU 1	MST 1	NAE 1
NAR 1	NDE 1	NRU 1	NSP C1
PVU 1	PVU 2	RRU 1	RSA 1
SAC 1	SAC 2	SAL 1	SMA 1
SNA 1	SPH 1	STU 1	TAQ 1
XBA 1	XHO 1	XHO 2	XMA 3
XMN 1			

10	20	30	40	50	60
GTTGTTGCTG	TGGCTGATAG	CCCCAGCAGG	GCCTGCACCT	GTGTCCCACC	CCACCCACAG
		SH			
		AA			
		UE			
		13			

70	80	90	100	110	120
ACGGCCTTCT	GCAATTCCGA	CCTCGTCATC	AGGGCCAAGT	TCGTGGGGAC	ACCAGAAGTC
H	T	M	SH		H
A	T	N	AA		I
E	H	L	UE		N
3	1	1	13		2

130	140	150	160	170	180
AACCAGACCA	CCTTATACCA	GCGTTATGAG	ATCAAGATGA	CCAAGATGTA	TAAAGGGTTC
		S			
		A			
		U			
		A			
190	200	210	220	230	240
CAAGCCTTAG	GGGATGCCGC	TGACATCCGG	TTCGTCTACA	CCCCCGCCAT	GGAGAGTGTC
SD	FS FN	F H	A	N	
AD	OF NS	O P	C	C	
UE	KA UP	K A	C	O	
11	11 12	1 2	1	1	
250	260	270	280	290	300
TGCGGATACT	TCCACAGGTC	CCACAACCGC	AGCGAGGAGT	TTCTCATTGC	TGGAAAAC TG
	A	B	M		P
	V	B	N		S
	A	V	L		T
	2	1	1		1
310	320	330	340	350	360
CAGGATGGAC	TCTTGACACAT	CACTACCTGC	AGTTTCGTGG	CTCCCTGGAA	CAGCCTGAGC
F H		P		B	D A D
O I		S		S	D L D
K N		T		T	E U E
1 1		1		1	1 1 1
370	380	390	400	410	420
TTAGCTCAGC	GCCGGGGCTT	CACCAAGACC	TACACTGTTG	GCTGTGAGGA	ATGCACAGTG
A D HH	N H			M	
L D AH	C P			N	
U E EA	I H			L	
1 1 21	1 1			1	
430	440	450	460	470	480
TTTCCCTGTT	TATCCATCCC	CTGCAAAC TG	CAGAGTGGCA	CTCATTGCTT	GTGGACGGAC
	F	P			A
	O	S			V
	K	T			A
	1	1			2
490	500	510	520	530	540
CAGCTCCTCC	AAGGCTCTGA	AAAGGGCTTC	CAGTCCCGTC	ACCTTGCC TG	CCTGCCTCGG
A M			H		MA
L N			P		NV
U L			H		LA
1 1			1		11

550	560	570	580	590	600
GAGCCAGGGC	TGTGCACCTG	GCAGTCCCTG	CGGTCCCAGA	TAGCCTGAAT	CCTGCCCCGGA
B	H	B	A	H	NH
S	G	S	V	I	CP
T	I	T	A	N	IA
1	1	1	2	1	12

610	620	630	640	650	660
GTGGAAGCTG	AAGCCTGCAC	AGTGTCCACC	CTGTTCCAC	TCCCATCTTT	CTTCCGGACA
A				M	H
L				B	P
U				O	A
1				2	2

670	680	690	700
ATGAAATAAA	GAGTTACCAC	CCAGCAAAAA	AAAAAAGGAA
			TTC
			E
			C
			O
			1

EXAMPLE 2-EXPRESSION OF COLLAGENASE INHIBITOR IN *E. COLI*

[0102] In this Example, a preferred method of coupling a preferred portable DNA sequence to the 5' end of the cloned cDNA is set forth. This involves making a nucleolytic cleavage at a specified point within the coding sequence and reconstructing the desired portion of the coding sequence by means of synthetic oligonucleotides in a manner that allows its excision and recombination (*i.e.*, by incorporating useful restriction sites).

[0103] Trimming the 5' end of the coding region will be accomplished by synthesizing both strands of the DNA extending from the *Tth111I* site in the 5' direction and ending in a *Bam*HI overhang. This synthetic oligonucleotide, referred to as FIBAC A, has the following features:

- (1) Codon selection has been biased toward those most frequently found in the genes of highly expressed bacterial proteins;

- (2) A methionine codon from which to initiate translation has been provided immediately upstream from the cysteine which begins the coding region of human processed FIBAC;
- (3) The spacing of the *Bam*HI site to the methionine codon is such that when cloned into pUC8, the coding region of FIBAC will be in-frame with the 5' end of the beta-galactosidase gene;
- (4) An in-frame stop codon and Shine Dalgarno sequence are also presented. Translation of this frame for the amino terminal portion of the beta-galactosidase is terminated at the TAA codon, and translation of FIBAC should be initiated at the following ATG;
- (5) Codons have been selected to create a *Hgi*AI site beginning with the G in the FIBAC initiation codon; and
- (6) There is a *Pvu*I site separated by one base from the 3' end of the *Bam*HI sequence.

[0104] The structure of FIBAC A is

GA TCC GCG ATC GGA GTG TAA GAA ATG TGC ACT
G CGC TAG CCT CAC ATT CTT TAC ACG TGA

TGC GTT CCG CCG CAT CCG CAG ACT GCT TTC
ACG CAA GGC GGC GTA GGC GTC TGA CGA AAG

TGC AAC TCT GAC C
ACG TTG AGA CTG CA

[0105] FIBAC A is synthesized using the ABI DNA synthesizer (Foster City, California) as a series of four component oligonucleotides.

[0106] Component oligonucleotide FA1 is:
GATCC GCGAT CGGAG TG TAA GAAAT GTGCA CTTGC

SVB C13)

[0107] Component oligonucleotide FA2 is:

GGAACG CAAGT GCACA TTTCT TACAC TCCGA TCGCG

SVB C14)

[0108] Component oligonucleotide FA3 is:

GTTC CGCCG CATCC GCAGA CTGCT TTCTG CAACT CTGAC C

SVB C15)

[0109] Component oligonucleotide FA4 is:

AGGTC AGACT TGCAG AAAGC AGTCT GCGGA TGCGG C

[0110] The remainder of the coding portion of the FIBAC gene is isolated as the 3' *Tth111I* to *EcoRI* fragment generated by a double digest of pUC9-F5/237P10 with these enzymes.

[0111] A synthetic linker is made to couple the 3' end of the *Tth111I* to *EcoRI* fragment to a *SalI* site. These oligonucleotides will be designed to recreate the *SalI* site and destroy the *EcoRI* site. The linker is comprised of the oligonucleotides linker A1 and linker A2.

SVB C16)

[0112] Linker A1 is: AATTGGCAG

SVB C17)

[0113] Linker A2 is: TCGACTGCC

[0114] These oligonucleotides and oligonucleotides FA1-FA4 are kinased separately and annealed in equal molar ratios with the *Tth111I* to *EcoRI* 3' end of the cDNA and *BamHI/SalI* cut mp19RF DNA. The ligated DNA is used to transfect JM105. Plaques are picked by their color in the presence of IPTG and X-gal and by hybridization to oligonucleotide FA2. Several positive plaques are to be sequenced. Those containing the designed sequence are subcloned into *BamHI/SalI* digested pUC8. Translation of the FIBAC gene in this construct is coupled to translation initiated for beta-galactosidase. This expression vector is referred to as pUC8-Fic.

[0115] Coupling translation of FIBAC to translation initiated for other highly expressed proteins is similarly arranged. For example, a portion of the *OmpA* gene

which contains the Shine-Dalgarno and initiator methionine sequences has been synthesized. This sequence encodes the entire signal peptide of OmpA protein and had convenient restriction sites, including those for *EcoRI*, *EcoRV*, *PvuI*, and *StuI*.

[0116] The sequence of the sense strand is:

10 20 30 40 50 60
GAATTCGATA TCTCGTTGGA GATATTCATG ACGTATTTTG GATGATAACG AGGCGCAAAA
E T E F M H
C A C O N H
O Q O K L A
1 1 5 1 1 1

70 80 90 100 110
AATGAAAAAG ACAGCTATCG CGATCGCAGT GGCACCTGGCT GGTTCGCTA CCGTA
A NF PS
L RN VA
U UU UU
1 12 1A

120 130
GCGCA GGCCTCTGGT AAAAGCTT
H S H M HA
H T A N IL
A U E L NU
1 1 3 1 31

[0117] This sequence is hereinafter referred to as OmpA leader. Coupling the translation of FIBAC to OmpA is accomplished by cutting the pUC8-Fic with *PvuI* and *SaI* and isolating the coding region. This, together with the *EcoRI* to *PvuI* fragment isolated from OmpA leader, will be cloned into *EcoRI*/*SaI*-cut pUC8. As in the prior example, transcription is driven by the lac promoter and regulated by the lac I gene product at the lac operator. This FIBAC expression vector is referred to as pUC8-F/OmpAic.

[0118] To effect the translocation of FIBAC out of the inner cell membrane, an appropriate leader sequence is added to the amino terminus of FIBAC. The protein thus produced will be translocated and processed to yield the mature form.

[0119] To effect such a translocation, a FIBAC gene encoding the signal peptide of the *E. coli* OmpA protein continuous with the structural region of FIBAC is created. This particular FIBAC gene necessitates having in frame stop codons at the 5' end of the FIBAC coding region changed. To accomplish this, the portion of the 5' coding region from pUC8-Fic that extends from the *Hgi*AI site to the *Nco*I site is isolated. Upstream sequences are resynthesized as a linker having cohesive ends from *Bam*HI and *Hgi*AI and containing an internal *Stu*I site. This is synthesized as two oligonucleotides, linker B1 and linker B2.

SUB C19) [0120] Linker B1 is: GATCCCAGGCCTGCA

SUB C20) [0121] Linker B2 is: GGCCTGG

[0122] Linkers B1 and B2 are kinased separately and annealed in equal molar ratios with the *Hgi*AI to *Nco*I fragment described above and *Bam*HI/*Nco*I cut pUC8-Fic. The resulting construct has the coding sequence of FIBAC in frame with the translation of the amino terminus of beta-galactosidase. Translation of this sequence forms a fusion protein with FIBAC. This plasmid is referred to as pUC8-Ff.

[0123] Attaching the OmpA leader sequences to the coding region of FIBAC is accomplished by ligating *Eco*RI/*Stu*I cut pUC8-Ff with an excess of the purified *Eco*RI to *Stu*I fragment of OmpA leader. Following transformation, plasmids from several colonies will be characterized by hybridization. Those that have incorporated the OmpA leader fragment are characterized further to verify the structure. This plasmid, pUC8-F OmpA1, will direct the synthesis of a fusion protein beginning in the signal peptide of the *E. coli* OmpA protein and ending in human FIBAC. The signals present in the

OmpA portion of the protein effect the protein's export from the cytoplasm and appropriate cleavage from the primary structure of FIBAC.

[0124] If the efficiency of expression were to be compromised by the sequence of the leader peptide or its combination with FIBAC either at the protein or at the nucleic acid level, the gene could be altered to encode any of several known *E. coli* leader sequences.

[0125] Transcription of all of the genes discussed is effected by the lac promoter. As in the case of initiation sites for translation, the promoter and operator region of the gene may be interchanged. FIBAC may also be expressed from vectors incorporating the lambda P_L promoter and operator (O_L), and the hybrid promoter operator, Tac as described in Amann, E., Brosius, J., and Ptashne, M. Gene 25: 167-178 (1983), specifically incorporated herein by reference. Excision of those portions of the gene including ribosome binding site structural region and 3' non-translated sequences and insertion in alternate vectors containing the P_L or Tac promoter makes use of the unique restriction sites that flank these structures in pUC8-F/OmpAic and pUC8-F/OmpA1. Insertion of the *EcoRI* to *SalI* fragment from either into similarly digested plasmid pDP8 effects transcription of these genes directed by the lambda P_L promoter. Transcriptional regulation would be temperature sensitive by merit of the cl857 mutation harbored on this same plasmid.

[0126] Putting similar gene fragments into the transcription unit of the Tac promoter will be accomplished by first isolating the *EcoRV* to *SalI* fragment. This, together with the synthetic Tac promoter sequence which is flanked by *BamHI* and *PvuII* sites and which contains the lac operator will be inserted into the *BamHI* to *SalI*

sites of pBR322 or preferably derivatives. The derivatives in this case refer to constructs containing either the *lacI* gene or the *l^q* gene.

[0127] Expression of FIBAC in host microorganisms other than *Escherichia* is considered. Yeast and bacteria of the genera *Bacillus*, *Pseudomonas*, and *Clostridium* may each offer particular advantages. The processes outlined above could easily be adapted to others.

[0128] In general, expression vectors for any microorganism will embody features analogous to those which we have incorporated in the above mentioned vectors of *E. coli*. In some cases, it will be possible to simply move the specific gene constructs discussed above directly into a vector compatible with the new host. In others, it may be necessary or desirable to alter certain operational or structural elements of the gene.

EXAMPLE 3

[0129] The human collagenase inhibitor may be readily purified after expression in a variety of microbes. In each case, the spectrum of contaminant proteins will differ. Thus, appropriate purification steps will be selected from a variety of steps already known to give a good separation of the human collagenase inhibitor from other proteins and from other procedures which are likely to work.

[0130] If the inhibitor is not secreted from the microbes, it may form inclusion bodies inside the recombinant microbes. These bodies are separated from other proteins by differential centrifugation after disruption of the cells with a French Press. The insoluble inclusion bodies are solubilized in 6 M guanidine hydrochloride or 8 M urea, and the inhibitor protein is more completely solubilized by reaction of its cysteines with sodium sulfite. At any time subsequent to this step, the cysteines are converted

back to their reduced form with dithiothreitol. Once the inhibitor protein is solubilized from inclusion bodies, immunoaffinity chromatography using antibodies raised against the unfolded inhibitor are used for purification before refolding.

[0131] The inhibitor can be refolded according to the protocol mentioned in Example 6, *infra*. After refolding of the inhibitor, or if the inhibitor is secreted from the microbes, purification from other proteins is accomplished by a variety of methods. Initial steps include ultrafiltration through a 50 k Dalton cutoff membrane or ammonium sulfate fractionation. Other useful methods include, but are not limited to, ion-exchange chromatography, gel filtration, heparin-sepharose chromatography, reversed-phase chromatography, or zinc-chelate chromatography. All of these steps have been successfully used in purification protocols. Additional high resolution steps include hydrophobic interaction chromatography or immunoaffinity chromatography. After purification, the metalloproteinase inhibitor is preferably at least 90-95% pure.

EXAMPLE 4

Purification of Human Collagenase Inhibitor from Human Amniotic Fluid

[0132] Human amniotic fluid obtained from discarded amniocentesis samples was pooled and 6 liters were subjected to ultrafiltration through a 100 kD MW cutoff filter, obtained from Millipore Corporation, in a Millipore Pellicon Cassette System. The eluate was concentrated through a 10 kD cutoff filter, obtained from Millipore Corporation, then through an Amicon PM-10 membrane. Aliquots (10 ml) of concentrated amniotic fluid were eluted through a 2.5 x 100 cm column of Ultrogel AcA54, obtained from LKB Corporation, which was equilibrated with pH 7.6, 0.05 M HEPES, 1 M sodium chloride, 0.01 M calcium chloride, and 0.02% sodium azide (all chemicals were obtained from Sigma Chemical Company). Fractions containing the

inhibitor were collected and pooled, dialyzed against pH 7.5, 0.025 M Hepes buffer containing 0.01 M calcium chloride and 0.02% sodium azide, and loaded onto a 1.5 x 28 cm heparin-sepharose CL-6B (obtained from Pharmacia, Inc.) column equilibrated with the same buffer. This column was rinsed with 1 liter of the above buffer and eluted with a linear gradient of 0-0.3 M sodium chloride. The fractions from the largest peak of inhibitor activity, eluting at about 0.1-0.15 M sodium chloride, were pooled, concentrated to 1 ml, and loaded onto a Synchropak rp-8 reverse phase HPLC column equilibrated with 0.05% trifluoroacetic acid (Aldrich Chemical Company). The column was eluted with a linear gradient of 0-40% acetonitrile (J. T. Baker Chemical Company) at 1/2% per minute. All fractions were immediately dried in a Savant speed-vac concentrator to remove acetonitrile, and redissolved in pH 7.5, 0.1 M Hepes before assay. The inhibitor eluted between 32-38% acetonitrile. Fractions containing the inhibitor were pooled, and 100 µl aliquots were eluted over a Bio-Rad biosil-TSK 250 HPLC gel filtration column. The pooled peaks of inhibitor activity contained 0.1 mg of inhibitor, which was over 95% pure as judged by SDS-polyacrylamide gel electrophoresis.

EXAMPLE 5

Purification of Human Fibroblast Collagenase Inhibitor from Human Embryonic Skin Fibroblast Serum-Free Medium

[0133] Human embryonic skin fibroblasts were grown in serum-free tissue culture medium. Ten liters of this medium were collected, dialyzed against pH 7.5, 0.02 M Hepes buffer containing 0.02% sodium azide and 0.01 M calcium chloride, and applied to a 2.8 x 48 cm column of heparin-sepharose CL-6B (Pharmacia, Inc.) equilibrated with the same buffer. The column was rinsed with 2 liters of this buffer and

was then eluted with linear gradient of 0-0.3 M sodium chloride contained in this buffer. The fractions obtained were tested for the presence of inhibitor by their ability to inhibit human fibroblast collagenase. The fractions corresponding to the peak of activity were those obtained near 0.15 M sodium chloride. These fractions were concentrated to about 5 ml by ultrafiltration through an Amicon YM10 filter and the concentrate was applied in four separate runs to a 250 x 4.1 mm Synchropak rp-8 reverse phase HPLC column, equilibrated with 1% trifluoroacetic acid. The column was eluted with a 0-60% linear gradient of acetonitrile in 0.1% trifluoroacetic acid. The gradient was run at 1/2% acetonitrile per minute. The inhibitor eluted in two sharp peaks between 26-29% acetonitrile. All fractions were immediately dried in a Savant speed-vac concentrator, redissolved in pH 7.5, 0.1 M Hepes, and assayed. At least 1.2 mg of collagenase inhibitor was recovered, which was 90-95% pure. This material gives a single band when run on a 17.5% reducing SDS gel. After carboxymethylation of the cysteines and elution through the same rp-8 column under identical conditions, the inhibitor is suitably homogenous for protein sequencing.

EXAMPLE 6

[0134] It is contemplated that the human collagenase inhibitor can be readily refolded into its native structure from its denatured state after expression of its gene in a microbe and separation of the collagenase inhibitor from most of the other proteins produced by the microbe. By analogy to the conditions necessary for the refolding of other disulfide-containing proteins as set forth by Freedman, R. B. and Hillson, D. A., in "Formation of Disulfide Bonds," In: The Enzymology of Post-Translational Modification of Proteins, Vol. 1, R. B. Freedman and H. C. Hawkins, eds., pp. 158-207 (1980), specifically incorporated herein by reference, refolding of the human collagenase

inhibitor should occur in solutions with a pH of 8.0 or greater. At this pH, the cysteines of the protein are partially ionized, and this condition is necessary for the attainment of native disulfide bond pairings. The inhibitor concentration should be relatively low, less than 0.1 mg/ml, to minimize the formation of intermolecular disulfide-linked aggregates which will interfere with the refolding process.

[0135] Since the stability of the refolded (native) disulfide bonded structure relative to the unfolded (reduced) structure depends on both the solution oxidation-reduction potential and the concentrations of other redox-active molecules, it is contemplated that the redox potential should be buffered with a redox buffer giving a potential equivalent to a reduced: oxidized glutathione ratio of 10. The preferred concentration range of reduced glutathione would be 0.1-1.0 mM. At higher concentrations, mixed disulfides will form with protein, reducing the yield of the refolded (native) structure. The relative stabilities of the unfolded protein and the native structure, and thus the rate and yield of refolding, will also depend on other solution variables, such as the pH, temperature, type of hydrogen-ion buffer, ionic strength, and the presence or absence of particular anions or cations as discussed in Privalov, P. L., "Stability of Proteins, Small Globular Proteins," in Advances in Protein Chemistry, Vol. 33, pp. 167-236, (1979), specifically incorporated herein by reference. These conditions vary for every protein and can be determined experimentally. It is contemplated that addition of any molecule that strongly prefers to bind the native (as opposed to the unfolded) structure, and which can be readily separated afterwards from the native (refolded) protein, will increase not only the yield but the rate of re-folding. These molecules include monoclonal antibodies raised against the native structure, and

other proteins which tightly bind the native collagenase inhibitor, such as the mammalian enzymes collagenase or gelatinase.

EXAMPLE 7

[0136] The second preferred sequence as set forth herein, *i.e.*,

10 20 30 40 50 60
GGCCATCGCC GCAGATCCAG CGCCCAGAGA GACACCAGAG AACCCACCAT GGCCCCCTTT
H F XB HH N S
A N HI AH C A
E U ON EA O U
3 1 21 21 1 1

70 80 90 100 110 120
GACCCCTGGC TTCTGCATCC TGTGTGTGCT GTGGCTGATA GCCCCAGCAG GGCCTGCACC
B SF S H
S FO A A
T AK U E
1 11 1 3

130 140 150 160 170 180
TGTGTCCCAC CCCACCCACA GACGGCCTTC TGCAATTCCG ACCTCGTCAT CAGGGCCAAG
H T M SH
A T N AA
E H L UE
3 1 1 13

190 200 210 220 230 240
TTCGTGGGGA CACCAGAAGT CAACCAGACC ACOTTATACC AGCGTTATGA GATCAAGATG
H S
I A
N U
2 A

250 260 270 280 290 300
ACCAAGATGT ATAAAGGGTT CCAAGCCTTA GGGGATGCCG CTGACATCCG GTTCGTCTAC
SD FS FN F H A
AD OF NS O P C
UE KA UP K A C
11 11 12 1 2 1

310 320 330 340 350 360
ACCCCCGCCA TGGAGAGTGT CTGCGGATAC TTCCACAGGT CCCACAACCG CAGCGAGGAG
N A B M
C V B N
O A V L
1 2 1 1

205270-2528660

SVB (21)

370	380	390	400	410	420
TTTCTCATTG	CTGGAAACT	GCAGGATGGA	CTCTTGCACA	TCACTACCTG	CAGTTTCGTG
	P	F	H	P	
	S	O	I	S	
	T	K	N	T	
	1	1	1	1	

430
GCTCCCTGGA AC
B
S
T
1

has the following restriction sites:

	#	SITES	FRAGMENTS		FRAGMENTS	ENDS
ACC 1 (GTVWAC)	1	295	295	(68.3)	1	295
			137	(31.7)	295	432
AVA 2 (GGRCC)	1	338	338	(78.2)	1	338
			94	(21.8)	338	432
BBV 1 (GCTGC)	1	350	350	(81.0)	1	350
			82	(19.0)	350	432
BIN 1 (GGATC)	1	14	418	(96.8)	14	432
			14	(3.2)	1	14
BST N1 (CCRGG)	2	65	360	(83.3)	65	425
		425	65	(15.0)	1	65
			7	(1.6)	425	432

205270-45E4B660

	#	SITES	FRAGMENTS	FRAGMENTS	ENDS
DDE 1 (CTNAG)	1				
		267	267 (61.8)	1	267
			165 (38.2)	267	432
FNU4H 1 (GCNGC)	3				
		8	269 (62.3)	8	277
		277	82 (19.0)	350	432
		350	73 (16.9)	277	350
			8 (1.9)	1	8
	#	SITES	FRAGMENTS	FRAGMENTS	ENDS
FOK 1 (GGATG)	4				
		76	197 (45.6)	76	273
		273	99 (22.9)	285	384
		285	76 (17.6)	1	76
		384	48 (11.1)	384	432
			12 (2.8)	273	285
HAE 2 (PGCGCQ)	1				
		19	413 (95.6)	19	432
			19 (4.4)	1	19
HAE 3 (GGCC)	5				
		1	258 (59.7)	174	432
		51	60 (13.9)	51	111
		111	50 (11.6)	1	51
		144	33 (7.6)	111	144
		174	30 (6.9)	144	174
			1 (0.2)	1	1
HHA 1 (GCGC)	1				
		20	412 (95.4)	20	432
			20 (4.6)	1	20
HINC 2 (GTQPAC)	1				
		199	233 (53.9)	199	432
			199 (46.1)	1	199

	#	SITES	FRAGMENTS		FRAGMENTS	ENDS
HINF 1 (GATC)	1	389	389	(90.0)	1	389
			43	(10.0)	389	432
HPA 2 (CCGG)	1	288	288	(66.7)	1	288
			144	(33.3)	288	432
MNL 1 (CCTC)	2	162	193	(44.7)	162	355
		355	162	(37.5)	1	162
			77	(17.8)	355	432
MST 2 (CCTNAGG)	1	266	266	(61.6)	1	266
			166	(38.4)	266	432
NCO 1 (CCATGG)	2	47	261	(60.4)	47	308
		308	124	(28.7)	308	432
			47	(10.9)	1	47
NSP B2 (CVGCWG)	1	278	278	(64.4)	1	278
			154	(35.6)	278	432
PST 1 (CTGCAG)	2	379	379	(87.7)	1	379
		408	29	(6.7)	379	408
			24	(5.6)	408	432
SAU 1 (CCTNAGG)	1	266	266	(61.6)	1	266
			166	(38.4)	266	432

	#	SITES	FRAGMENTS		FRAGMENTS	ENDS
SAU 3A (GATC)	2					
		14	217	(50.2)	14	231
		231	201	(46.5)	231	432
			14	(3.2)	1	14
SAU96 1 (GGNCC)	4					
		51	165	(38.2)	173	338
		110	94	(21.8)	338	432
		173	63	(14.6)	110	173
		338	59	(13.7)	51	110
			51	(11.8)	1	51
SCR F1 (CCNGG)	2					
		65	360	(83.3)	65	425
		425	65	(15.0)	1	65
			7	(1.6)	425	432
SFA N1 (GATGC)	2					
		75	199	(46.1)	75	274
		274	158	(36.6)	274	432
			75	(17.4)	1	75
STY 1 (CCRRGG)	2					
		47	261	(60.4)	47	308
		308	124	(28.7)	308	432
			47	(10.9)	1	47
TTH111 I (GACNNNGTC)	1					
		160	272	(63.0)	160	432
			160	(37.0)	1	160
XHO 2 (PGATCQ)	1					
		13	419	(97.0)	13	432
			13	(3.0)	1	13

[0137] The following do not appear:

AAT 2	AFL 2	AFL 3	AHA 2
AHA 3	ALU 1	APA 1	ASU 2
AVA 1	AVA 3	AVR 2	BAL 1
BAM H1	BAN1	BAN 2	BCL 1
BGL 1	BGL 2	BSM 1	BSP 1286
nSSH 1	BST E2	CFR 1	CLA 1
ECO R1	ECO R5	FNUD 2	GDI 2
HAE 1	HGA 1	HGI A1	HGI CI
HGI D1	HGI J2	HIND 3	HPA 1
HPH 1	KPN 1	MBO 2	MLU 1
MST 1	NAE 1	NAR 1	NCI 1
NDE 1	NHE 1	NOT 1	NRU 1
NSP C1	PVU 1	PVU 2	RRU 1
RSA 1	SAC 1	SAC 2	SAL 1
SCA 1	SMA 1	SNA 1	SNA B1
SPE 1	SPH 1	SSP 1	STU 1
TAQ 1	XBA 1	XHO 1	XMA 3
XMN 1			

[0138] The salient features of this cDNA are:

1. The coding strand is presented in the 5' to 3' convention with the polyC tract at the 5' end.
2. If the first G in the sequence GGC CAT CGC CGC is considered as nucleotide 1, then an open reading frame exists from nucleotide 1 through nucleotide 432, which is the 3' end of this partial cDNA.
3. The first methionine in this reading frame is encoded by nucleotides 49 through 51 and represents the initiation site of translation.
4. The amino acid sequence prescribed by nucleotides 49 through 114 is not found in the primary structure of the mature protein, but it is the sequence of the leader peptide of human protein.
5. The sequence of nucleotides 82 through 432 is identical to the sequence of nucleotides numbered 1 through 351 in the insert from the first preferred sequence of Example 1.
6. The amino acid sequence of the mature protein displays two consensus sequences for sugar attachment. These sequences, -N-Q-T- prescribed by nucleotides 202 through 210 and -N-R-S- prescribed by nucleotides 346 through 354, are amino acid residues 30 through 32 and 78 through

80, respectively, in the mature protein. Both sites are glycosylated in the human inhibitor protein.

[0140] It will be apparent to those skilled in the art that various modifications and variations can be made in the processes and products of the present invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents.

205270" /SE/28660